List of Figures

| Figure 1-1 | Planning Area Chugach National Forest | |
|---------------|--|------|
| Figure 2-1 | Alternative development | 2-3 |
| Figure 2-2 | Graphic summary of the range of | |
| | alternatives, given as percent of total Forest | |
| | acreage by prescription category | 2-16 |
| Figure 2-3 | No Action Alternative area allocations, by | - 1- |
| | category | 2-18 |
| Figure 2-4 | Preferred Alternative area allocations, by | 0.40 |
| | category | |
| Figure 2-5 | Alternative A area allocations, by category | |
| Figure 2-6 | Alternative B area allocations, by category | |
| Figure 2-7 | Alternative C area allocations, by category | |
| Figure 2-8 | Alternative D area allocations, by category | |
| Figure 2-9 | Alternative E area allocations, by category | |
| Figure 2-10 | Alternative F area allocations, by category | 2-25 |
| Figure 2-11 | Bioenvironmental classes of the Chugach | |
| | National Forest with at least 12 percent of | |
| | area in Category 1 or 2 prescriptions by | |
| | alternative | 2-27 |
| Figure 2-12 | Percent of Kenai portion of Chugach | |
| | National Forest heavily infested by spruce | |
| | bark beetle (mapped as infested for 3+ | |
| | years) by prescription category and | |
| | alternative | 2-27 |
| Figure 2-13 | Percentage of coho and pink salmon | |
| _ | habitat by prescription category | 2-29 |
| Figure 2-14 | Potential stream class I, II, and III | |
| _ | restoration miles | 2-29 |
| Figure 2-15 | Forestwide distribution of ROS classes | 2-35 |
| Figure 2-16 | Forestwide dispersed recreation capacity | 2-36 |
| Figure 2-17 | Acres available for winter snowmachine | |
| J | activity Kenai Peninsula | 2-37 |
| Figure 2-18 | Acres available for winter helicopter | |
| 3 | activities Kenai Peninsula | |
| Figure 2-19 | Recommended and existing Research | |
| 1.9 | Natural Areas by alternative (acres) | 2-38 |
| Figure 2-20 | Recommended Wilderness by alternative | |
| g | (acres) | 2-39 |
| Figure 2-21 | Percent of Forest roadless by alternative | |
| | , - | _ |

| Figure 3-1 | Total short-term soil disturbance from |
|---------------------------------------|--|
| | recreation and timber roads and trails by |
| | geographic area3-20 |
| Figure 3-2 | Total long-term soil disturbance from |
| | recreation and timber roads and trails by |
| | geographic area3-21 |
| Figure 3-3a | Ecological regions of the Chugach National |
| | Forest (CNF lands only)3-51 |
| Figure 3-3b | Ecological sections of the Chugach |
| ga. 0 0 0 0 | National Forest (CNF lands only)3-52 |
| Figure 3-4a | Vascular plant species richness and |
| i iguic o +u | community richness3-55 |
| Figure 3-4b | Vascular plant community richness by |
| i iguic 3-45 | formation class (i.e., level 1 of Viereck et al. |
| | 1992), and community richness3-55 |
| Figure 2.4c | , · |
| Figure 3-4c | Vascular plant species richness by |
| | formation class and geographic area (i.e., |
| | Kenai Peninsula, Prince William Sound and |
| Fig 0 F- | Copper River Delta)3-56 |
| Figure 3-5a | Area distribution of forested cover types on |
| | the Kenai Peninsula portion of the Chugach |
| | National Forest in the 1975 baseline and |
| | estimated for the years 1875, 2000, 2050, |
| | and 2100 using the Kenai Forest |
| | Succession Model3-59 |
| Figure 3-5b | Area distribution of tree age classes on the |
| | Kenai Peninsula portion of the Chugach |
| | National Forest in the 1975 baseline and |
| | estimated for the years 1875, 2000, 2050, |
| | and 2100 using the Kenai Forest |
| | Succession Model3-60 |
| Figure 3-6 | Pollen percentages over approximately the |
| | last 2,500 years from sites at Tern Lake, |
| | Circle Lake and Hidden Lake3-65 |
| Figure 3-7a | Predicted changes in tree basal area in |
| - | response to climatic change for a site in the |
| | Kenai Mountains at 300 meter elevation, on |
| | a westerly aspect with a slope of 25 percent3-66 |
| Figure 3-7b | Regional Landscape Area3-79 |
| Figure 3-8a | Proportion of land cover class by |
| | prescription category by alternative3-83 |
| Figure 3-8b | Proportions of vegetative cover types by |
| J 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 | prescription category, by alternative (does |
| | not include ANILCA additions)3-88 |
| | 110t 11101000 / 1111EO/ t ddditio110/0-00 |

| Figure 3-8c | Proportion of forest structural classes by | |
|---------------|--|-------|
| | alternative, all forest types (does not | |
| | include ANILCA additions) | 3-89 |
| Figure 3-9a | Bioenvironmental classes with at least 12 | |
| | percent of area in Category 1 or 2 | |
| | prescriptions by alternative | 3-91 |
| Figure 3-9b | Area of bioenvironmental classes with less | |
| | than 12 percent of area in Category 1 or 2 | |
| | prescriptions by alternative | 3-92 |
| Figure 3-10 | Comparison of species richness for | |
| • | summer, migration, and winter by | |
| | alternative | 3-99 |
| Figure 3-11 | Percentage of coho and pink salmon | |
| 9 | habitat by prescription category | 3-111 |
| Figure 3-12 | Percentage of watershed associations | |
| | within Category 1 and 2 prescriptions | 3-113 |
| Figure 3-13 | Percentage of watershed associations | |
| ga. e e . e | harvested under high market conditions | 3-120 |
| Figure 3-14 | Estimated miles of class I, II, and III | |
| 1 19410 0 1 1 | restoration by alternative | 3-126 |
| Figure 3-15a | Fire history of National Forest lands on the | |
| rigare o roa | Kenai Peninsula from 1914 to 1997 as | |
| | indicated by the number of fires and acres | |
| | burned by decade | 3-129 |
| Figure 3-15b | Area by fuels model class on the Kenai | 5-129 |
| rigule 3-13b | | |
| | Peninsula portion of the Chugach National Forest | 3-132 |
| Figure 2 16 | | 3-132 |
| Figure 3-16 | Spruce bark beetle infestation acreage on | |
| | the Kenai Peninsula portion of the Chugach | |
| | National Forest for the years 1957, 1969, | 0.400 |
| Г: 0. 47- | and 1976 through 1998 | 3-139 |
| Figure 3-17a | Cumulative acres of proposed timber | 0.440 |
| E' - 0 47b | harvest and stand improvement - decade 1 | 3-143 |
| Figure 3-17b | Cumulative acres of vegetation treated by | |
| | fuels, wildlife, and forest restoration | 0.444 |
| = 0.40 | programs - decade 1 | 3-144 |
| Figure 3-18 | Acres of forested land on the Chugach | |
| | National Forest in Category 1 prescriptions | |
| | by alternative | 3-147 |
| Figure 3-19 | Cumulative acres of treated and untreated | |
| | forestland - decade 1 | 3-148 |
| Figure 3-20 | Cumulative acres of treated and untreated | |
| | forestland - decade 5 | 3-149 |
| Figure 3-21 | Forested land distribution by geographic | |
| | landscape (acres), Chugach National | |
| | Forest | 3-153 |

| Figure 3-22 | Structural stages of forested lands on Chugach National Forest | 3_155 |
|-------------|--|-------|
| Figure 3-23 | Percent of available timberlands by forest | 5-100 |
| rigure 3-23 | type and structural stage on the Chugach | |
| | National Forest | 3 155 |
| Eiguro 2 24 | Acres by structural stage for available | 5-133 |
| Figure 3-24 | | |
| | timberlands on the Chugach National Forest | 3-156 |
| Figure 2 0F | | 3-130 |
| Figure 3-25 | Forest density classes on the Chugach | 0.457 |
| Figure 2.00 | National Forest | 3-157 |
| Figure 3-26 | Vegetation management on the Chugach | 0.450 |
| F: 0.07 | National Forest by activity, 1974-1999 | 3-158 |
| Figure 3-27 | Distribution of potential Aphragmus | |
| | eschscholtzianus habitat on the Chugach | |
| | National Forest by prescription category | 0.400 |
| E: 0.00 | and alternative | 3-192 |
| Figure 3-28 | Distribution of potential Arnica lessingii spp. | |
| | norbergii habitat on the Chugach National | |
| | Forest by prescription category and | |
| | | 3-193 |
| Figure 3-29 | Distribution of potential Carex lenticularis | |
| | var. dolia habitat on the Chugach National | |
| | Forest by prescription category and | |
| | alternative | 3-194 |
| Figure 3-30 | Distribution of potential Draba kananaskis | |
| | habitat on the Chugach National Forest by | |
| | prescription category and alternative | 3-195 |
| Figure 3-31 | Distribution of potential Isoetes truncata | |
| | habitat on the Chugach National Forest by | |
| | prescription category and alternative | 3-196 |
| Figure 3-32 | Distribution of potential Ligusticum calderi | |
| | habitat on the Chugach National Forest by | |
| | prescription category and alternative | 3-197 |
| Figure 3-33 | Distribution of potential Papaver | |
| J | alboroseum habitat on the Chugach | |
| | National Forest by prescription category | |
| | , , , , , , , , , , , , , , , , , , , | 3-198 |
| Figure 3-34 | Distribution of potential Puccinellia glabra | |
| J | habitat on the Chugach National Forest by | |
| | prescription category and alternative | 3-199 |
| Figure 3-35 | Distribution of potential Romanzoffia | |
| J | unalaschcensis habitat on the Chugach | |
| | National Forest by prescription category | |
| | and alternative | 3-200 |
| | | |

| Figure 3-36 | Distribution of potential Stellaria ruscifolia | |
|----------------|--|-------|
| | ssp. aleutica habitat on the Chugach | |
| | National Forest by prescription category | |
| | and alternative | 3-201 |
| Figure 3-37 | Distribution of potential black oystercatcher | |
| · · | habitat on the Chugach National Forest by | |
| | prescription category (21,500 acres) | 3-234 |
| Figure 3-38 | Distribution of potential brown bear habitat | |
| 5 | on the Chugach National Forest by | |
| | prescription category (1,828,190 acres) | 3-237 |
| Figure 3-39 | Distribution of potential dusky Canada | |
| 9 | goose habitat on the Chugach National | |
| | Forest by prescription category (541,750 | |
| | acres) | 3-238 |
| Figure 3-40 | Percent of prescription categories on the | 200 |
| rigaro o ro | Kenai Peninsula that may affect moose | |
| | winter habitat (482,100 acres) | 3-240 |
| Figure 3-41 | Percent of prescription categories on the | 2 10 |
| riguic o +1 | Copper River Delta that may affect moose | |
| | winter habitat (256,860 acres) | 3-241 |
| Figure 3-42 | Comparison of H.S.I. values when 3,300 | 5-241 |
| riguic 3-42 | acres per year of moose winter habitat are | |
| | burned in the next 5 years and no | |
| | management (natural succession) | 3-243 |
| Figure 3-43 | Comparison of H.S.I. values when 1,200 | 5-245 |
| rigule 3-43 | acres per year of moose winter habitat are | |
| | burned in the next 5 years and no | |
| | | 3-243 |
| Figure 2 44 | management (natural succession) | 3-243 |
| Figure 3-44 | Distribution of potential mountain goat | |
| | habitat on the Chugach National Forest by | 2 245 |
| Figure 0, 45 a | prescription category (1,336,300 acres) | 3-245 |
| Figure 3-45a | Distribution of potential gray wolf habitat on | |
| | the Chugach National Forest by | 0.047 |
| E: 0.4E! | prescription category (2,742,210 acres) | 3-247 |
| Figure 3-45b | Distribution of potential gray wolf habitat on | |
| | the Copper River Delta by prescription | |
| | category (783,740 acres) | 3-247 |
| Figure 3-46 | Distribution of potential lynx habitat on the | |
| | Chugach National Forest by prescription | |
| | category (2,665,700 acres) | 3-248 |
| Figure 3-47 | Distribution of potential marbled murrelet | |
| | habitat on the Chugach National Forest by | |
| | prescription category (1,455,520 acres) | 3-250 |
| Figure 3-48 | Distribution of potential river otter habitat on | |
| | the Chugach National Forest by | |
| | prescription category (1,981,100 acres) | 3-251 |

| Figure 3-49 Distribution of potential Sitka black-tailed deer habitat on the Chugach National | | |
|---|--|-------|
| | Forest by prescription category (1,239,620 | |
| | acres) | 3-253 |
| Figure 3-50 | Distribution of potential Townsend's warbler | |
| | habitat on the Chugach National Forest by | |
| | prescription category (734,280 acres) | 3-254 |
| Figure 3-51 | Distribution of potential wolverine habitat on | |
| | the Chugach National Forest by | |
| | prescription category (3,153,870 acres) | 3-255 |
| Figure 3-52 | Distribution of potential trumpeter swan | |
| | habitat on the Chugach National Forest by | |
| | prescription category (628,410 acres) | |
| Figure 3-53 | Kenai Peninsula geographic area | |
| Figure 3-53a | Campground occupancy – Kenai Peninsula | |
| Figure 3-53b | Cabin occupancy – Kenai Peninsula | |
| Figure 3-53c | Inventoried ROS classes – Kenai Peninsula | 3-309 |
| Figure 3-53d | Recreation capacities of dispersed areas – | |
| | Kenai Peninsula | 3-309 |
| Figure 3-54 | Prince William Sound geographic area | 3-311 |
| Figure 3-54a | Cabin occupancy – Prince William Sound | 3-314 |
| Figure 3-54b | Inventoried ROS classes – Prince William | |
| | Sound | 3-315 |
| Figure 3-54c | Recreation capacities of dispersed areas – | |
| | Prince William Sound | 3-315 |
| Figure 3-55 | Copper River Delta geographic area | |
| Figure 3-55a | Cabin occupancy – Copper River Delta | 3-319 |
| Figure 3-55b | Inventoried ROS classes – Copper River | |
| | Delta | 3-320 |
| Figure 3-55c | Recreation capacities of dispersed areas – | |
| | Copper River Delta | 3-320 |
| Figure 3-56a | Numbers of U.S. adults participating in | |
| | selected outdoor recreation activities in | |
| | 1983, 1995, and 2000 | 3-325 |
| Figure 3-56b | Median age of U.S. population from 1940 to | |
| | 2050 | 3-327 |
| Figure 3-56c | Percent change in U.S. adult participation in | |
| | selected recreation activities, 1995 to 2010 | 3-328 |
| Figure 3-57a | Participation rates by Alaskan adult | |
| | population and U.S. adult population in | |
| | outdoor recreation activities in 1995 | 3-329 |
| Figure 3-57b | Number of Alaskan adults participating in | |
| | outdoor recreation activities in 1995 | 3-330 |
| Figure 3-57c | Alaska statewide outdoor recreation | |
| | participation estimates, 2000-2020, using | |
| | the 1995 NSRE database | 3-331 |

| Figure 3-57d | Increase in number of adult Alaskans | |
|--------------|---|-------|
| | participating in recreation activities, from 2000 to 2020 | 3-331 |
| Figure 3-57e | Average number of times an Alaskan adult | |
| J | annually participates in recreation activities | 3-333 |
| Figure 3-57f | Projected increases in total times Alaskans | |
| J | participate in recreation activities, 2000 to | |
| | 2020 | 3-334 |
| Figure 3-58a | Total visitors entering Alaska during May | |
| _ | through September, 1989 to 1999 | 3-335 |
| Figure 3-58b | Rate of growth in total summer visitors to | |
| J | Alaska, 1989 to 1999 | 3-336 |
| Figure 3-58c | Projections of participation by U.S. adults | |
| _ | (excluding Alaskans) in wildlife-related | |
| | recreation activities in Alaska | 3-337 |
| Figure 3-59a | Percentage participation in recreation | |
| J | activities on the Chugach National Forest, | |
| | based on Recreation Survey '95 | 3-339 |
| Figure 3-59b | Participation rates of visitors and residents | |
| J | in recreation activities on the Chugach | |
| | National Forest, summer 1995 | 3-340 |
| Figure 3-59c | Total visits to the Chugach National Forest | |
| J | to participate in recreation activities, based | |
| | on 1998 INFRA data | 3-342 |
| Figure 3-60a | Primitive ROS by alternative – Kenai | |
| _ | Peninsula | 3-351 |
| Figure 3-60b | Semi-primitive ROS by alternative – Kenai | |
| | Peninsula | 3-351 |
| Figure 3-60c | Roaded ROS by alternative – Kenai | |
| _ | Peninsula | 3-352 |
| Figure 3-60d | ROS settings by alternative – Kenai | |
| _ | Peninsula | 3-353 |
| Figure 3-60e | Projected cabin capacity vs. projected cabin | |
| | use – Kenai Peninsula | 3-355 |
| Figure 3-60f | Projected camping capacity vs. projected | |
| _ | camping use – Kenai Peninsula | 3-356 |
| Figure 3-60g | Developed recreation capacity vs. projected | |
| | use – Kenai Peninsula | 3-357 |
| Figure 3-60h | Dispersed recreation capacity – Kenai | |
| | Peninsula | 3-358 |
| Figure 3-60i | Acres available for winter snowmachine | |
| | activity – Kenai Peninsula | 3-359 |
| Figure 3-60j | Acres available for winter helicopter | |
| | activities – Kenai Peninsula | 3-360 |
| Figure 3-61a | Primitive ROS by alternative – Prince | |
| | William Sound | 3-361 |

| Figure 3-61b | Semi-primitive ROS by alternative – Prince William Sound | 3-362 |
|---------------|--|--------|
| Figure 3-61c | Roaded ROS by alternative – Prince | |
| | William Sound | 3-362 |
| Figure 3-61d | ROS setting by alternative – Prince William | 0.000 |
| Figure 3-61e | Sound Projected cabin capacity vs. projected cabin | 3-363 |
| rigule 3-01e | use – Prince William Sound | 3-365 |
| Figure 3-61f | Developed recreation capacity vs. projected | |
| J | use - Prince William Sound | 3-366 |
| Figure 3-61g | Dispersed recreation capacity – Prince | |
| | William Sound | 3-366 |
| Figure 3-62a | Primitive ROS by alternative – Copper | 0.000 |
| Fig 2 COb | River Delta | 3-368 |
| Figure 3-62b | Semi-primitive ROS by alternative – Copper River Delta | 3-368 |
| Figure 3-62c | Roaded ROS by alternative – Copper River | 5-500 |
| 1 19410 0 020 | Delta | 3-369 |
| Figure 3-62d | ROS setting by alternative – Copper River | |
| 3 | Delta | 3-370 |
| Figure 3-62e | Projected cabin capacity vs. projected cabin | |
| | use – Copper River Delta | 3-372 |
| Figure 3-62f | Projected camping capacity vs. projected | |
| F: 0.00 | camping use – Copper River Delta | 3-372 |
| Figure 3-62g | Developed recreation capacity vs. projected | 2 272 |
| Figure 3-63 | use – Copper River Delta Dispersed recreation capacity – Copper | 3-373 |
| i igule 3-03 | River Delta | 3-374 |
| Figure 3-64 | Subsistence use by six rural communities in | 0 01 4 |
| 9000 | or adjacent to the Chugach National Forest | 3-381 |
| Figure 3-65 | Percentage use of wild resources by | |
| _ | community | 3-384 |
| Figure 3-66 | Location of the eight eligible RNAs, | |
| | including the existing Green Island RNA | 0.000 |
| F' 0 07 | within the Chugach National Forest | 3-393 |
| Figure 3-67 | Bioenvironmental class representation | |
| | within the eligible network of eight RNAs sorted by total area of each | |
| | bioenvironmental class | 3-394 |
| Figure 3-68 | Existing scenic integrity | |
| Figure 3-69 | Scenic attractiveness | |
| Figure 3-70 | Landscape visibility analysis by geographic | |
| - | area | |
| Figure 3-71 | Scenic classes | 3-426 |
| Figure 3-72 | Mapped scenic integrity objectives by | |
| | alternative | 3-430 |

| Figure 3-73 | Inventoried Roadless Areas map | 3-454 |
|--------------|---|-------|
| Figure 3-74 | Historical timber harvest on the Chugach | |
| | National Forest 1909-1999 | 3-469 |
| Figure 3-75 | Historical timber harvest on the Chugach | |
| | National Forest, 1909-1999, by decade | 3-470 |
| Figure 3-76 | Southcentral Alaska timber harvest by | |
| | ownership, 1986-1996 | 3-473 |
| Figure 3-77 | Sawlog harvest volume | 3-473 |
| Figure 3-78 | Utility harvest volume | 3-473 |
| Figure 3-79 | Population growth, 1979-1997, indexed to | |
| | 1979 | 3-511 |
| Figure 3-80 | Total employment trends, 1979-1997, | |
| | indexed to 1979 | 3-512 |
| Figure 3-81 | Total personal income trends, 1979-1997, | |
| _ | for the United States, Alaska, and | |
| | Southcentral Alaska – in 1999 dollars, | |
| | indexed to 1979 | 3-515 |
| Figure 3-82 | Indexed per capita income for the United | |
| _ | States, Alaska, and Southcentral Alaska, | |
| | 1979-1997 in 1999 dollars | 3-516 |
| Figure 3-83 | Municipality of Anchorage employment by | |
| | sector highlighting forest resource related | |
| | sectors 1996 | 3-520 |
| Figure 3-84 | Kenai and Soldotna employment by sector | |
| | highlighting forest resource related sectors, | |
| | 1996 | 3-521 |
| Figure 3-85 | Other communities within or near the | |
| | Chugach National Forest boundary | |
| | employment by sector highlighting forest | |
| | resource related sectors, 1996 | 3-522 |
| Figure 3-85a | Real income in the commercial fishing | |
| | sector, 1979-98 | 3-524 |
| Figure 3-85b | Index of income in the visitor industry | |
| _ | (adjusted for inflation), 1979-98 | 3-526 |
| Figure 3-85c | Major components of visitor industry by | |
| _ | share of total visitor industry income, 1998 | 3-527 |
| Figure 3-86 | Chugach National Forest total revenue in | |
| _ | 1999 dollars, fiscal years 1980-1998 | 3-529 |
| Figure 3-87 | Preference for the amount of designated | |
| · · | Wilderness | 3-554 |
| Figure 3-88 | Preference for the amount of designated | |
| · · | Wild and Scenic Rivers | 3-555 |
| Figure 3-89 | Preference for the amount of timber harvest | |
| Figure 3-90 | Acceptable conditions for timber harvest | |
| Figure 3-91 | Preference for the amount of new roads | |
| Figure 3-92 | Acceptable purposes for road construction | |
| _ | | |

List of Figures

| Figure 3-93 Preference for open areas for snowm use | |
|---|-------|
| Figure 3-94 Preference for open areas for off high | |
| vehicles | • |
| Figure 3-95 Preference for management response | |
| the Whittier road | 3-562 |
| Figure B-1 Process for identification of lands suit | able |
| for timber production | B-11 |